

Zeta functions of K3 surfaces over finite fields

Kiran S. Kedlaya

Department of Mathematics, University of California San Diego

kedlaya@ucsd.edu

These slides can be downloaded from <https://kskedlaya.org/slides/>.

Arithmetic, Geometry and Computations on K3 Surfaces

ICERM, Providence, RI

September 17, 2026

Supported by  (grant DMS-2401536) and  (Warschawski Professorship).

I acknowledge that my workplace occupies unceded ancestral land of the Kumeyaay Nation.



Contents

- 1 Prelude on zeta functions
- 2 Structure of the zeta function of a K3 surface
- 3 Positivity considerations
- 4 Computational aspects
- 5 Lattice-theoretic considerations
- 6 Conclusion

Hasse–Weil zeta functions

For X a scheme of finite type over $\mathbb{Z}$, the **zeta function** is the complex analytic function defined for $\text{Real}(s) \gg 0$ by the Euler product

$$\zeta(X, s) := \prod_{x \in X^\circ} (1 - \#\kappa(x)^{-s})^{-1}$$

where X° runs over closed points of the scheme X and $\kappa(x)$ is the residue field of x .

For example, for $X = \text{Spec } \mathbb{Z}$ this gives the Riemann zeta function. More generally, for K a number field and $\mathfrak{o}_K$ the ring of integers, taking $X = \text{Spec } \mathfrak{o}_K$ gives the Dedekind zeta function of K .

Hasse–Weil zeta functions

For X a scheme of finite type over $\mathbb{Z}$, the **zeta function** is the complex analytic function defined for $\text{Real}(s) \gg 0$ by the Euler product

$$\zeta(X, s) := \prod_{x \in X^\circ} (1 - \#\kappa(x)^{-s})^{-1}$$

where X° runs over closed points of the scheme X and $\kappa(x)$ is the residue field of x .

For example, for $X = \text{Spec } \mathbb{Z}$ this gives the Riemann zeta function. More generally, for K a number field and $\mathfrak{o}_K$ the ring of integers, taking $X = \text{Spec } \mathfrak{o}_K$ gives the Dedekind zeta function of K .

Zeta functions of varieties over a finite field

Hereafter, fix a prime power $q = p^f$ and let $\mathbb{F}_q$ be the finite field of order q . For X an algebraic variety (or any scheme of finite type) over $\mathbb{F}_q$, we have

$$\zeta(X, s) = Z(X, q^{-s})$$

for the power series $Z(X, T)$ over $\mathbb{Z}$ given by

$$Z(X, T) = \exp \left(\sum_{n=1}^{\infty} \frac{T^n}{n} \#X(\mathbb{F}_{q^n}) \right).$$

This power series always represents a **rational function**. For example, for X a genus 1 curve,

$$Z(X, T) = \frac{1 - aT + qT^2}{(1 - T)(1 - qT)} \text{ where } a := q + 1 - \#X(\mathbb{F}_q) \text{ satisfies } |a| \leq 2\sqrt{q} \text{ (Hasse).}$$

Equivalently, the numerator factors over $\mathbb{C}$ as $(1 - \alpha T)(1 - \beta T)$ with $|\alpha| = |\beta| = q^{1/2}$.

Zeta functions of varieties over a finite field

Hereafter, fix a prime power $q = p^f$ and let $\mathbb{F}_q$ be the finite field of order q . For X an algebraic variety (or any scheme of finite type) over $\mathbb{F}_q$, we have

$$\zeta(X, s) = Z(X, q^{-s})$$

for the power series $Z(X, T)$ over $\mathbb{Z}$ given by

$$Z(X, T) = \exp \left(\sum_{n=1}^{\infty} \frac{T^n}{n} \#X(\mathbb{F}_{q^n}) \right).$$

This power series always represents a **rational function**. For example, for X a genus 1 curve,

$$Z(X, T) = \frac{1 - aT + qT^2}{(1 - T)(1 - qT)} \text{ where } a := q + 1 - \#X(\mathbb{F}_q) \text{ satisfies } |a| \leq 2\sqrt{q} \text{ (Hasse).}$$

Equivalently, the numerator factors over $\mathbb{C}$ as $(1 - \alpha T)(1 - \beta T)$ with $|\alpha| = |\beta| = q^{1/2}$.

Zeta functions of varieties over a finite field

Hereafter, fix a prime power $q = p^f$ and let $\mathbb{F}_q$ be the finite field of order q . For X an algebraic variety (or any scheme of finite type) over $\mathbb{F}_q$, we have

$$\zeta(X, s) = Z(X, q^{-s})$$

for the power series $Z(X, T)$ over $\mathbb{Z}$ given by

$$Z(X, T) = \exp \left(\sum_{n=1}^{\infty} \frac{T^n}{n} \#X(\mathbb{F}_{q^n}) \right).$$

This power series always represents a **rational function**. For example, for X a genus 1 curve,

$$Z(X, T) = \frac{1 - aT + qT^2}{(1 - T)(1 - qT)} \text{ where } a := q + 1 - \#X(\mathbb{F}_q) \text{ satisfies } |a| \leq 2\sqrt{q} \text{ (Hasse).}$$

Equivalently, the numerator factors over $\mathbb{C}$ as $(1 - \alpha T)(1 - \beta T)$ with $|\alpha| = |\beta| = q^{1/2}$.

Zeta functions of abelian varieties

For X an abelian variety of dimension g over $\mathbb{F}_q$, Weil showed that¹

$$Z(X, T) = \prod_{i=0}^{2g} (\wedge^i L_X)(T)^{(-1)^{i+1}} = \frac{(\wedge^1 L_X)(T)(\wedge^3 L_X)(T) \cdots}{(\wedge^0 L_X)(T)(\wedge^2 L_X)(T) \cdots}$$

for a certain polynomial $L_X(T)$ of degree $2g$ which satisfies

$$L_X(0) = 1, \quad q^g L_X(q^{-1}T) = T^{2g} L_X(qT^{-1})$$

and the reciprocal roots $\alpha_1, \dots, \alpha_{2g}$ in $\mathbb{C}$ satisfy $|\alpha_i| = q^{1/2}$. More concretely,

$$\#X(\mathbb{F}_{q^n}) = \prod_{i=1}^{2g} (1 - \alpha_i^n) \quad (n = 1, 2, \dots).$$

¹Notation gloss: If $F: V \rightarrow V$ satisfies $\det(1 - FT, V) = L_X(T)$, then $\det(1 - FT, \wedge^i V) = (\wedge^i L_X)(T)$.

The inverse problem for abelian varieties

By Tate's theorem, the zeta function is a **complete isogeny invariant** for abelian varieties over $\mathbb{F}_q$. One can refine this to get a complete **isomorphism** invariant; more on this later.

Moreover, it is known exactly which polynomials $L_X(T)$ can occur in the zeta function of some abelian variety X (Honda–Tate, Milne–Waterhouse): the obstruction cannot occur for $q = p$, is rare for $q \neq p$, and is (easily) explicitly computable. Using this, one can tabulate isogeny classes of abelian varieties; see [LMFDB](#).

The inverse problem for abelian varieties

By Tate's theorem, the zeta function is a **complete isogeny invariant** for abelian varieties over $\mathbb{F}_q$. One can refine this to get a complete **isomorphism** invariant; more on this later.

Moreover, it is known exactly which polynomials $L_X(T)$ can occur in the zeta function of some abelian variety X (Honda–Tate, Milne–Waterhouse): the obstruction cannot occur for $q = p$, is rare for $q \neq p$, and is (easily) explicitly computable. Using this, one can tabulate isogeny classes of abelian varieties; see [LMFDB](#).

Contents

- 1 Prelude on zeta functions
- 2 Structure of the zeta function of a K3 surface
- 3 Positivity considerations
- 4 Computational aspects
- 5 Lattice-theoretic considerations
- 6 Conclusion

Goals of this talk

By a **K3 surface** over a field k of any characteristic, I mean a smooth, projective, geometrically irreducible surface X which is simply connected² with trivial canonical bundle.

This talk will address two general questions about K3 surfaces (and related varieties).

- What can we say about zeta functions of K3 surfaces over finite fields?
- What does this tell us about the geometry and arithmetic of K3 surfaces?

Importantly, neither of these questions is entirely concentrated in the setting of working over a finite field! On one hand, some of the information we get about zeta functions over finite fields comes from working over $\overline{\mathbb{F}}_q$ or even lifting to characteristic 0. On the other hand, the information we get is also relevant in characteristic 0 (cf. Edgar's minicourse).

²That is, every finite étale covering of the base extension $X_{\bar{k}}$ splits.

Goals of this talk

By a **K3 surface** over a field k of any characteristic, I mean a smooth, projective, geometrically irreducible surface X which is simply connected² with trivial canonical bundle.

This talk will address two general questions about K3 surfaces (and related varieties).

- What can we say about zeta functions of K3 surfaces over finite fields?
- What does this tell us about the geometry and arithmetic of K3 surfaces?

Importantly, neither of these questions is entirely concentrated in the setting of working over a finite field! On one hand, some of the information we get about zeta functions over finite fields comes from working over $\overline{\mathbb{F}}_q$ or even lifting to characteristic 0. On the other hand, the information we get is also relevant in characteristic 0 (cf. Edgar's minicourse).

²That is, every finite étale covering of the base extension $X_{\bar{k}}$ splits.

Goals of this talk

By a **K3 surface** over a field k of any characteristic, I mean a smooth, projective, geometrically irreducible surface X which is simply connected² with trivial canonical bundle.

This talk will address two general questions about K3 surfaces (and related varieties).

- What can we say about zeta functions of K3 surfaces over finite fields?
- What does this tell us about the geometry and arithmetic of K3 surfaces?

Importantly, neither of these questions is entirely concentrated in the setting of working over a finite field! On one hand, some of the information we get about zeta functions over finite fields comes from working over $\overline{\mathbb{F}}_q$ or even lifting to characteristic 0. On the other hand, the information we get is also relevant in characteristic 0 (cf. Edgar's minicourse).

²That is, every finite étale covering of the base extension $X_{\bar{k}}$ splits.

Goals of this talk

By a **K3 surface** over a field k of any characteristic, I mean a smooth, projective, geometrically irreducible surface X which is simply connected² with trivial canonical bundle.

This talk will address two general questions about K3 surfaces (and related varieties).

- What can we say about zeta functions of K3 surfaces over finite fields?
- What does this tell us about the geometry and arithmetic of K3 surfaces?

Importantly, neither of these questions is entirely concentrated in the setting of working over a finite field! On one hand, some of the information we get about zeta functions over finite fields comes from working over $\overline{\mathbb{F}}_q$ or even lifting to characteristic 0. On the other hand, the information we get is also relevant in characteristic 0 (cf. Edgar's minicourse).

²That is, every finite étale covering of the base extension $X_{\bar{k}}$ splits.

Goals of this talk

By a **K3 surface** over a field k of any characteristic, I mean a smooth, projective, geometrically irreducible surface X which is simply connected² with trivial canonical bundle.

This talk will address two general questions about K3 surfaces (and related varieties).

- What can we say about zeta functions of K3 surfaces over finite fields?
- What does this tell us about the geometry and arithmetic of K3 surfaces?

Importantly, neither of these questions is entirely concentrated in the setting of working over a finite field! On one hand, some of the information we get about zeta functions over finite fields comes from working over $\overline{\mathbb{F}}_q$ or even lifting to characteristic 0. On the other hand, the information we get is also relevant in characteristic 0 (cf. Edgar's minicourse).

²That is, every finite étale covering of the base extension $X_{\overline{k}}$ splits.

Motivation: Kuga–Satake

Over $\mathbb{C}$, the Kuga–Satake construction presents the Hodge structure of a K3 surface using the tensor square of the Hodge structure of some (large) abelian variety. Using the language of **Shimura varieties**, this construction can be described in a way that makes sense over finite fields, even³ in characteristic 2.

For example, this relationship was used by Deligne to establish the properties of zeta functions of K3 surfaces **before** he had proved the Weil conjectures in general. See next slide for a precise statement.

It is also used to prove the Tate conjecture for $H^2(X)$ and $H^2(X \times_{\mathbb{F}_q} X)$ (Ito–Ito–Koshikawa + much earlier work). We will (silently) use many consequences of this later.

This intuition also helps us formulate other questions and conjectures...

³It will be a running motif in this talk that “2 is the oddest of all characteristics.”

Motivation: Kuga–Satake

Over $\mathbb{C}$, the Kuga–Satake construction presents the Hodge structure of a K3 surface using the tensor square of the Hodge structure of some (large) abelian variety. Using the language of **Shimura varieties**, this construction can be described in a way that makes sense over finite fields, even³ in characteristic 2.

For example, this relationship was used by Deligne to establish the properties of zeta functions of K3 surfaces **before** he had proved the Weil conjectures in general. See next slide for a precise statement.

It is also used to prove the Tate conjecture for $H^2(X)$ and $H^2(X \times_{\mathbb{F}_q} X)$ (Ito–Ito–Koshikawa + much earlier work). We will (silently) use many consequences of this later.

This intuition also helps us formulate other questions and conjectures...

³It will be a running motif in this talk that “2 is the oddest of all characteristics.”

Motivation: Kuga–Satake

Over $\mathbb{C}$, the Kuga–Satake construction presents the Hodge structure of a K3 surface using the tensor square of the Hodge structure of some (large) abelian variety. Using the language of **Shimura varieties**, this construction can be described in a way that makes sense over finite fields, even³ in characteristic 2.

For example, this relationship was used by Deligne to establish the properties of zeta functions of K3 surfaces **before** he had proved the Weil conjectures in general. See next slide for a precise statement.

It is also used to prove the Tate conjecture for $H^2(X)$ and $H^2(X \times_{\mathbb{F}_q} X)$ (Ito–Ito–Koshikawa + much earlier work). We will (silently) use many consequences of this later.

This intuition also helps us formulate other questions and conjectures...

³It will be a running motif in this talk that “2 is the oddest of all characteristics.”

Motivation: Kuga–Satake

Over $\mathbb{C}$, the Kuga–Satake construction presents the Hodge structure of a K3 surface using the tensor square of the Hodge structure of some (large) abelian variety. Using the language of **Shimura varieties**, this construction can be described in a way that makes sense over finite fields, even³ in characteristic 2.

For example, this relationship was used by Deligne to establish the properties of zeta functions of K3 surfaces **before** he had proved the Weil conjectures in general. See next slide for a precise statement.

It is also used to prove the Tate conjecture for $H^2(X)$ and $H^2(X \times_{\mathbb{F}_q} X)$ (Ito–Ito–Koshikawa + much earlier work). We will (silently) use many consequences of this later.

This intuition also helps us formulate other questions and conjectures...

³It will be a running motif in this talk that “2 is the oddest of all characteristics.”

Zeta functions of K3 surfaces (after Deligne)

For X a K3 surface over $\mathbb{F}_q$,

$$Z(X, T) = (1 - T)^{-1}(1 - q^2 T)^{-1} q Q(qT)^{-1}$$

where the polynomial $Q(T)$ satisfies $Q(T) \in q + T\mathbb{Z}[T]$, $\deg(Q) = 22$, $Q(1) = 0$, and

$$Q(T) = \prod_{i=1}^{22} (1 - \alpha_i T), \quad \alpha_i \in \mathbb{C}, \quad |\alpha_i| = 1.$$

The polynomial $Q(T)$ is reciprocal up to sign:

$$T^{22} Q(T^{-1}) = (-1)^{\text{ord}_{T=-1}(Q)} Q(T).$$

Question (“Honda–Tate problem”)

What other restrictions are there on Q ?

Zeta functions of K3 surfaces (after Deligne)

For X a K3 surface over $\mathbb{F}_q$,

$$Z(X, T) = (1 - T)^{-1}(1 - q^2 T)^{-1} q Q(qT)^{-1}$$

where the polynomial $Q(T)$ satisfies $Q(T) \in q + T\mathbb{Z}[T]$, $\deg(Q) = 22$, $Q(1) = 0$, and

$$Q(T) = \prod_{i=1}^{22} (1 - \alpha_i T), \quad \alpha_i \in \mathbb{C}, \quad |\alpha_i| = 1.$$

The polynomial $Q(T)$ is reciprocal up to sign:

$$T^{22} Q(T^{-1}) = (-1)^{\text{ord}_{T=-1}(Q)} Q(T).$$

Question (“Honda–Tate problem”)

What other restrictions are there on Q ?

Zeta functions of K3 surfaces (after Deligne)

For X a K3 surface over $\mathbb{F}_q$,

$$Z(X, T) = (1 - T)^{-1}(1 - q^2 T)^{-1} q Q(qT)^{-1}$$

where the polynomial $Q(T)$ satisfies $Q(T) \in q + T\mathbb{Z}[T]$, $\deg(Q) = 22$, $Q(1) = 0$, and

$$Q(T) = \prod_{i=1}^{22} (1 - \alpha_i T), \quad \alpha_i \in \mathbb{C}, \quad |\alpha_i| = 1.$$

The polynomial $Q(T)$ is reciprocal up to sign:

$$T^{22} Q(T^{-1}) = (-1)^{\text{ord}_{T=-1}(Q)} Q(T).$$

Question (“Honda–Tate problem”)

What other restrictions are there on Q ?

Zeta functions of K3 surfaces (after Deligne)

For X a K3 surface over $\mathbb{F}_q$,

$$Z(X, T) = (1 - T)^{-1}(1 - q^2 T)^{-1} q Q(qT)^{-1}$$

where the polynomial $Q(T)$ satisfies $Q(T) \in q + T\mathbb{Z}[T]$, $\deg(Q) = 22$, $Q(1) = 0$, and

$$Q(T) = \prod_{i=1}^{22} (1 - \alpha_i T), \quad \alpha_i \in \mathbb{C}, \quad |\alpha_i| = 1.$$

The polynomial $Q(T)$ is reciprocal up to sign:

$$T^{22} Q(T^{-1}) = (-1)^{\text{ord}_{T=-1}(Q)} Q(T).$$

Question (“Honda–Tate problem”)

What other restrictions are there on Q ?

Factoring the zeta function

There is a unique factorization of polynomials $Q(T) = Q_{\text{alg}}(T)Q_{\text{trans}}(T)$ where

- $Q_{\text{alg}}(0) = 1$ and **every** root of Q_{alg} is a root of unity;
- $Q_{\text{trans}}(0) = 1$ and **no** root of Q_{trans} is a root of unity.

The degree of $Q_{\text{alg}}(T)$ is the **geometric** Picard rank $\bar{\rho} = \text{rank Pic}(X_{\overline{\mathbb{F}}_q})$, which is always even. By contrast, the Picard rank $\rho = \text{rank Pic}(X)$ equals $\text{ord}_{T=1}(Q_{\text{alg}}) = \text{ord}_{T=1}(Q)$ which has no parity constraint.

Note that $\bar{\rho} = 22$ is possible even though this cannot occur over $\mathbb{C}$! In this case we say X is **supersingular** (e.g., the Fermat quartic when $q \equiv 3 \pmod{4}$). However, Artin showed that $\rho \neq 22$ if $q = p^f$ with f odd.

Question

For X supersingular, are there any other constraints on Q_{alg} in terms of q ? Special case: for every $m \in \{-20, \dots, 22\}$, can we have $\#X(\mathbb{F}_q) = q^2 + mq + 1$ for suitable q, X ?

Factoring the zeta function

There is a unique factorization of polynomials $Q(T) = Q_{\text{alg}}(T)Q_{\text{trans}}(T)$ where

- $Q_{\text{alg}}(0) = 1$ and **every** root of Q_{alg} is a root of unity;
- $Q_{\text{trans}}(0) = 1$ and **no** root of Q_{trans} is a root of unity.

The degree of $Q_{\text{alg}}(T)$ is the **geometric** Picard rank $\bar{\rho} = \text{rank Pic}(X_{\overline{\mathbb{F}}_q})$, which is always even. By contrast, the Picard rank $\rho = \text{rank Pic}(X)$ equals $\text{ord}_{T=1}(Q_{\text{alg}}) = \text{ord}_{T=1}(Q)$ which has no parity constraint.

Note that $\bar{\rho} = 22$ is possible even though this cannot occur over $\mathbb{C}$! In this case we say X is **supersingular** (e.g., the Fermat quartic when $q \equiv 3 \pmod{4}$). However, Artin showed that $\rho \neq 22$ if $q = p^f$ with f odd.

Question

For X supersingular, are there any other constraints on Q_{alg} in terms of q ? Special case: for every $m \in \{-20, \dots, 22\}$, can we have $\#X(\mathbb{F}_q) = q^2 + mq + 1$ for suitable q, X ?

Factoring the zeta function

There is a unique factorization of polynomials $Q(T) = Q_{\text{alg}}(T)Q_{\text{trans}}(T)$ where

- $Q_{\text{alg}}(0) = 1$ and **every** root of Q_{alg} is a root of unity;
- $Q_{\text{trans}}(0) = 1$ and **no** root of Q_{trans} is a root of unity.

The degree of $Q_{\text{alg}}(T)$ is the **geometric** Picard rank $\bar{\rho} = \text{rank Pic}(X_{\overline{\mathbb{F}}_q})$, which is always even. By contrast, the Picard rank $\rho = \text{rank Pic}(X)$ equals $\text{ord}_{T=1}(Q_{\text{alg}}) = \text{ord}_{T=1}(Q)$ which has no parity constraint.

Note that $\bar{\rho} = 22$ is possible even though this cannot occur over $\mathbb{C}$! In this case we say X is **supersingular** (e.g., the Fermat quartic when $q \equiv 3 \pmod{4}$). However, Artin showed that $\rho \neq 22$ if $q = p^f$ with f odd.

Question

For X supersingular, are there any other constraints on Q_{alg} in terms of q ? Special case: for every $m \in \{-20, \dots, 22\}$, can we have $\#X(\mathbb{F}_q) = q^2 + mq + 1$ for suitable q, X ?

Factoring the zeta function

There is a unique factorization of polynomials $Q(T) = Q_{\text{alg}}(T)Q_{\text{trans}}(T)$ where

- $Q_{\text{alg}}(0) = 1$ and **every** root of Q_{alg} is a root of unity;
- $Q_{\text{trans}}(0) = 1$ and **no** root of Q_{trans} is a root of unity.

The degree of $Q_{\text{alg}}(T)$ is the **geometric** Picard rank $\bar{\rho} = \text{rank Pic}(X_{\overline{\mathbb{F}}_q})$, which is always even. By contrast, the Picard rank $\rho = \text{rank Pic}(X)$ equals $\text{ord}_{T=1}(Q_{\text{alg}}) = \text{ord}_{T=1}(Q)$ which has no parity constraint.

Note that $\bar{\rho} = 22$ is possible even though this cannot occur over $\mathbb{C}$! In this case we say X is **supersingular** (e.g., the Fermat quartic when $q \equiv 3 \pmod{4}$). However, Artin showed that $\rho \neq 22$ if $q = p^f$ with f odd.

Question

For X supersingular, are there any other constraints on Q_{alg} in terms of q ? Special case: for every $m \in \{-20, \dots, 22\}$, can we have $\#X(\mathbb{F}_q) = q^2 + mq + 1$ for suitable q, X ?

Factoring the zeta function

There is a unique factorization of polynomials $Q(T) = Q_{\text{alg}}(T)Q_{\text{trans}}(T)$ where

- $Q_{\text{alg}}(0) = 1$ and **every** root of Q_{alg} is a root of unity;
- $Q_{\text{trans}}(0) = 1$ and **no** root of Q_{trans} is a root of unity.

The degree of $Q_{\text{alg}}(T)$ is the **geometric** Picard rank $\bar{\rho} = \text{rank Pic}(X_{\overline{\mathbb{F}}_q})$, which is always even. By contrast, the Picard rank $\rho = \text{rank Pic}(X)$ equals $\text{ord}_{T=1}(Q_{\text{alg}}) = \text{ord}_{T=1}(Q)$ which has no parity constraint.

Note that $\bar{\rho} = 22$ is possible even though this cannot occur over $\mathbb{C}$! In this case we say X is **supersingular** (e.g., the Fermat quartic when $q \equiv 3 \pmod{4}$). However, Artin showed that $\rho \neq 22$ if $q = p^f$ with f odd.

Question

For X supersingular, are there any other constraints on Q_{alg} in terms of q ? Special case: for every $m \in \{-20, \dots, 22\}$, can we have $\#X(\mathbb{F}_q) = q^2 + mq + 1$ for suitable q, X ?

Contents

- 1 Prelude on zeta functions
- 2 Structure of the zeta function of a K3 surface
- 3 Positivity considerations**
- 4 Computational aspects
- 5 Lattice-theoretic considerations
- 6 Conclusion

Lower bounds on point counts

Let X be a K3 surface over $\mathbb{F}_q$. By Deligne's theorem and the triangle inequality

$$\#X(\mathbb{F}_q) = q^2 + 1 + q \sum_{i=1}^{22} \alpha_i \geq q^2 + q + 1 - 21q = q^2 - 20q + 1.$$

In particular, if $q \geq 23$ then $X(\mathbb{F}_q) \neq \emptyset$.

Question

For which q does there exist a K3 surface X over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$?

Such examples are known for $q = 2, 3, 4, 5$. One example for $q = 5$ is the Fermat quartic.

Theorem (Bernasconi–Filipazzi, 2025)

Let $q = p^f$ be a prime power with $p > 5$ for which there exists no K3 surface X over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$ (e.g., $q = p \geq 23$). Then every smooth, projective, geometrically integral 3-fold over $\mathbb{F}_q$ with nef anticanonical class and negative Kodaira dimension has a rational point.

Lower bounds on point counts

Let X be a K3 surface over $\mathbb{F}_q$. By Deligne's theorem and the triangle inequality

$$\#X(\mathbb{F}_q) = q^2 + 1 + q \sum_{i=1}^{22} \alpha_i \geq q^2 + q + 1 - 21q = q^2 - 20q + 1.$$

In particular, if $q \geq 23$ then $X(\mathbb{F}_q) \neq \emptyset$.

Question

For which q does there exist a K3 surface X over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$?

Such examples are known for $q = 2, 3, 4, 5$. One example for $q = 5$ is the Fermat quartic.

Theorem (Bernasconi–Filipazzi, 2025)

Let $q = p^f$ be a prime power with $p > 5$ for which there exists no K3 surface X over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$ (e.g., $q = p \geq 23$). Then every smooth, projective, geometrically integral 3-fold over $\mathbb{F}_q$ with nef anticanonical class and negative Kodaira dimension has a rational point.

Lower bounds on point counts

Let X be a K3 surface over $\mathbb{F}_q$. By Deligne's theorem and the triangle inequality

$$\#X(\mathbb{F}_q) = q^2 + 1 + q \sum_{i=1}^{22} \alpha_i \geq q^2 + q + 1 - 21q = q^2 - 20q + 1.$$

In particular, if $q \geq 23$ then $X(\mathbb{F}_q) \neq \emptyset$.

Question

For which q does there exist a K3 surface X over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$?

Such examples are known for $q = 2, 3, 4, 5$. One example for $q = 5$ is the Fermat quartic.

Theorem (Bernasconi–Filipazzi, 2025)

Let $q = p^f$ be a prime power with $p > 5$ for which there exists no K3 surface X over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$ (e.g., $q = p \geq 23$). Then every smooth, projective, geometrically integral 3-fold over $\mathbb{F}_q$ with nef anticanonical class and negative Kodaira dimension has a rational point.

Lower bounds on point counts

Let X be a K3 surface over $\mathbb{F}_q$. By Deligne's theorem and the triangle inequality

$$\#X(\mathbb{F}_q) = q^2 + 1 + q \sum_{i=1}^{22} \alpha_i \geq q^2 + q + 1 - 21q = q^2 - 20q + 1.$$

In particular, if $q \geq 23$ then $X(\mathbb{F}_q) \neq \emptyset$.

Question

For which q does there exist a K3 surface X over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$?

Such examples are known for $q = 2, 3, 4, 5$. One example for $q = 5$ is the Fermat quartic.

Theorem (Bernasconi–Filipazzi, 2025)

Let $q = p^f$ be a prime power with $p > 5$ for which there exists no K3 surface X over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$ (e.g., $q = p \geq 23$). Then every smooth, projective, geometrically integral 3-fold over $\mathbb{F}_q$ with nef anticanonical class and negative Kodaira dimension has a rational point.

Positivity constraints

Turning this around, the condition $\#X(\mathbb{F}_q) \geq 0$ imposes a nontrivial constraint on the zeta functions for $q \leq 19$. Some other nontrivial constraints of this type are the following.

- $\#X(\mathbb{F}_{q^2}) \geq \#X(\mathbb{F}_q)$ (for $q = 2, 3, 4$)
- $\#X(\mathbb{F}_{q^3}) \geq \#X(\mathbb{F}_q)$ (for $q = 2$).
- $\#X(\mathbb{F}_{q^4}) \geq \#X(\mathbb{F}_{q^2})$ (for $q = 2$).

Aside: for K3 surfaces, the inverse problem for zeta functions is solved “up to base extension” if $p > 5$ (Taelman, Ito). Any attempt to eliminate the base extension would have to account for these positivity constraints.⁴

⁴These constraints would disappear if one allowed **noncommutative** K3 surfaces, for which the trace-theoretic point counts are not enumerative. It may be easier to prove an analogue of Honda–Tate in this setting.

Positivity constraints

Turning this around, the condition $\#X(\mathbb{F}_q) \geq 0$ imposes a nontrivial constraint on the zeta functions for $q \leq 19$. Some other nontrivial constraints of this type are the following.

- $\#X(\mathbb{F}_{q^2}) \geq \#X(\mathbb{F}_q)$ (for $q = 2, 3, 4$)
- $\#X(\mathbb{F}_{q^3}) \geq \#X(\mathbb{F}_q)$ (for $q = 2$).
- $\#X(\mathbb{F}_{q^4}) \geq \#X(\mathbb{F}_{q^2})$ (for $q = 2$).

Aside: for K3 surfaces, the inverse problem for zeta functions is solved “up to base extension” if $p > 5$ (Taelman, Ito). Any attempt to eliminate the base extension would have to account for these positivity constraints.⁴

⁴These constraints would disappear if one allowed **noncommutative** K3 surfaces, for which the trace-theoretic point counts are not enumerative. It may be easier to prove an analogue of Honda–Tate in this setting.

Positivity constraints

Turning this around, the condition $\#X(\mathbb{F}_q) \geq 0$ imposes a nontrivial constraint on the zeta functions for $q \leq 19$. Some other nontrivial constraints of this type are the following.

- $\#X(\mathbb{F}_{q^2}) \geq \#X(\mathbb{F}_q)$ (for $q = 2, 3, 4$)
- $\#X(\mathbb{F}_{q^3}) \geq \#X(\mathbb{F}_q)$ (for $q = 2$).
- $\#X(\mathbb{F}_{q^4}) \geq \#X(\mathbb{F}_{q^2})$ (for $q = 2$).

Aside: for K3 surfaces, the inverse problem for zeta functions is solved “up to base extension” if $p > 5$ (Taelman, Ito). Any attempt to eliminate the base extension would have to account for these positivity constraints.⁴

⁴These constraints would disappear if one allowed **noncommutative** K3 surfaces, for which the trace-theoretic point counts are not enumerative. It may be easier to prove an analogue of Honda–Tate in this setting.

Positivity constraints

Turning this around, the condition $\#X(\mathbb{F}_q) \geq 0$ imposes a nontrivial constraint on the zeta functions for $q \leq 19$. Some other nontrivial constraints of this type are the following.

- $\#X(\mathbb{F}_{q^2}) \geq \#X(\mathbb{F}_q)$ (for $q = 2, 3, 4$)
- $\#X(\mathbb{F}_{q^3}) \geq \#X(\mathbb{F}_q)$ (for $q = 2$).
- $\#X(\mathbb{F}_{q^4}) \geq \#X(\mathbb{F}_{q^2})$ (for $q = 2$).

Aside: for K3 surfaces, the inverse problem for zeta functions is solved “up to base extension” if $p > 5$ (Taelman, Ito). Any attempt to eliminate the base extension would have to account for these positivity constraints.⁴

⁴These constraints would disappear if one allowed **noncommutative** K3 surfaces, for which the trace-theoretic point counts are not enumerative. It may be easier to prove an analogue of Honda–Tate in this setting.

Positivity constraints

Turning this around, the condition $\#X(\mathbb{F}_q) \geq 0$ imposes a nontrivial constraint on the zeta functions for $q \leq 19$. Some other nontrivial constraints of this type are the following.

- $\#X(\mathbb{F}_{q^2}) \geq \#X(\mathbb{F}_q)$ (for $q = 2, 3, 4$)
- $\#X(\mathbb{F}_{q^3}) \geq \#X(\mathbb{F}_q)$ (for $q = 2$).
- $\#X(\mathbb{F}_{q^4}) \geq \#X(\mathbb{F}_{q^2})$ (for $q = 2$).

Aside: for K3 surfaces, the inverse problem for zeta functions is solved “up to base extension” if $p > 5$ (Taelman, Ito). Any attempt to eliminate the base extension would have to account for these positivity constraints.⁴

⁴These constraints would disappear if one allowed **noncommutative** K3 surfaces, for which the trace-theoretic point counts are not enumerative. It may be easier to prove an analogue of Honda–Tate in this setting.

A sharper Weil bound for Q_{trans}

One obtains better bounds on the contribution of Q_{trans} to $\#X(\mathbb{F}_{q^n})$ using the fact that for every cyclotomic polynomial Φ_i (of degree $\varphi(i)$),

$$\varphi(i) \log q + \sum_{\alpha: Q_{\text{trans}}(\alpha)=0} \log |\Phi_i(\alpha)| = \log |\text{Res}(Q_{\text{trans}}, \Phi_i)| \geq 0$$

plus a little bit of linear programming (and some exhaustive searches).

Theorem (K, 2026+)

Let X be a K3 surface over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$. If $q = 7, 8$, then $\bar{\rho} \geq 4$; if $q = 9$, then $\bar{\rho} \geq 6$; if $q = 11$, then $\bar{\rho} \geq 8$; if $q = 13$, then $\bar{\rho} \geq 10$; if $q = 16, 17$, then $\bar{\rho} \geq 16$; if $q = 19$, then $\bar{\rho} \geq 18$.

This means that one can potentially rule out the larger primes by analyzing a small number of options for $Q(T)$. (Note: if $\bar{\rho} = 22$ then $\#X(\mathbb{F}_q) \equiv 1 \pmod{q}$.)

A sharper Weil bound for Q_{trans}

One obtains better bounds on the contribution of Q_{trans} to $\#X(\mathbb{F}_{q^n})$ using the fact that for every cyclotomic polynomial Φ_i (of degree $\varphi(i)$),

$$\varphi(i) \log q + \sum_{\alpha: Q_{\text{trans}}(\alpha)=0} \log |\Phi_i(\alpha)| = \log |\text{Res}(Q_{\text{trans}}, \Phi_i)| \geq 0$$

plus a little bit of linear programming (and some exhaustive searches).

Theorem (K, 2026+)

Let X be a K3 surface over $\mathbb{F}_q$ with $X(\mathbb{F}_q) = \emptyset$. If $q = 7, 8$, then $\bar{\rho} \geq 4$; if $q = 9$, then $\bar{\rho} \geq 6$; if $q = 11$, then $\bar{\rho} \geq 8$; if $q = 13$, then $\bar{\rho} \geq 10$; if $q = 16, 17$, then $\bar{\rho} \geq 16$; if $q = 19$, then $\bar{\rho} \geq 18$.

This means that one can potentially rule out the larger primes by analyzing a small number of options for $Q(T)$. (Note: if $\bar{\rho} = 22$ then $\#X(\mathbb{F}_q) \equiv 1 \pmod{q}$.)

... and related varieties

From the zeta function $Z(X, T)$, one can compute the zeta functions of Hilbert schemes associated to X . This gives rise to additional positivity considerations.

Theorem (Debarre–Laface–Roulleau + ϵ)

For $q \neq 3$, any smooth cubic fourfold over $\mathbb{F}_q$ contains an $\mathbb{F}_q$ -rational line.

For $q \geq 5$, expressing the point count on the Fano variety of lines $F(X)$ in terms of $Z(X, T)$ gives a positive lower bound. For $q = 2, 4$, one instead shows that

$$\#F(X)(\mathbb{F}_q) \equiv 1 + m + m^2 \pmod{q}, \quad m := \text{Trace}(\text{Frob}_q, H^2(F(X), \mathbb{Q}_\ell)) \in \mathbb{Z}$$

and this cannot be 0 when q is a power of a prime $p \equiv 2 \pmod{3}$,

Since the only missing case is $q = 3$, and the moduli space of cubic fourfolds has roughly $3^{19} \approx 10^9$ points over $\mathbb{F}_3$, it is feasible to fill this gap using a census.

... and related varieties

From the zeta function $Z(X, T)$, one can compute the zeta functions of Hilbert schemes associated to X . This gives rise to additional positivity considerations.

Theorem (Debarre–Laface–Roulleau + ϵ)

For $q \neq 3$, any smooth cubic fourfold over $\mathbb{F}_q$ contains an $\mathbb{F}_q$ -rational line.

For $q \geq 5$, expressing the point count on the Fano variety of lines $F(X)$ in terms of $Z(X, T)$ gives a positive lower bound. For $q = 2, 4$, one instead shows that

$$\#F(X)(\mathbb{F}_q) \equiv 1 + m + m^2 \pmod{q}, \quad m := \text{Trace}(\text{Frob}_q, H^2(F(X), \mathbb{Q}_\ell)) \in \mathbb{Z}$$

and this cannot be 0 when q is a power of a prime $p \equiv 2 \pmod{3}$,

Since the only missing case is $q = 3$, and the moduli space of cubic fourfolds has roughly $3^{19} \approx 10^9$ points over $\mathbb{F}_3$, it is feasible to fill this gap using a census.

... and related varieties

From the zeta function $Z(X, T)$, one can compute the zeta functions of Hilbert schemes associated to X . This gives rise to additional positivity considerations.

Theorem (Debarre–Laface–Roulleau + ϵ)

For $q \neq 3$, any smooth cubic fourfold over $\mathbb{F}_q$ contains an $\mathbb{F}_q$ -rational line.

For $q \geq 5$, expressing the point count on the Fano variety of lines $F(X)$ in terms of $Z(X, T)$ gives a positive lower bound. For $q = 2, 4$, one instead shows that

$$\#F(X)(\mathbb{F}_q) \equiv 1 + m + m^2 \pmod{q}, \quad m := \text{Trace}(\text{Frob}_q, H^2(F(X), \mathbb{Q}_\ell)) \in \mathbb{Z}$$

and this cannot be 0 when q is a power of a prime $p \equiv 2 \pmod{3}$,

Since the only missing case is $q = 3$, and the moduli space of cubic fourfolds has roughly $3^{19} \approx 10^9$ points over $\mathbb{F}_3$, it is feasible to fill this gap using a census.

... and related varieties

From the zeta function $Z(X, T)$, one can compute the zeta functions of Hilbert schemes associated to X . This gives rise to additional positivity considerations.

Theorem (Debarre–Laface–Roulleau + ϵ)

For $q \neq 3$, any smooth cubic fourfold over $\mathbb{F}_q$ contains an $\mathbb{F}_q$ -rational line.

For $q \geq 5$, expressing the point count on the Fano variety of lines $F(X)$ in terms of $Z(X, T)$ gives a positive lower bound. For $q = 2, 4$, one instead shows that

$$\#F(X)(\mathbb{F}_q) \equiv 1 + m + m^2 \pmod{q}, \quad m := \text{Trace}(\text{Frob}_q, H^2(F(X), \mathbb{Q}_\ell)) \in \mathbb{Z}$$

and this cannot be 0 when q is a power of a prime $p \equiv 2 \pmod{3}$,

Since the only missing case is $q = 3$, and the moduli space of cubic fourfolds has roughly $3^{19} \approx 10^9$ points over $\mathbb{F}_3$, it is feasible to fill this gap using a census.

Contents

- 1 Prelude on zeta functions
- 2 Structure of the zeta function of a K3 surface
- 3 Positivity considerations
- 4 Computational aspects**
- 5 Lattice-theoretic considerations
- 6 Conclusion

Exhausting over Weil polynomials

SageMath includes code (by me) to exhaust over Weil polynomials satisfying various constraints. This was used to build the tables of isogeny classes of abelian varieties over finite fields in [LMFDB](#), but you can also use it directly.

```
sage: P.<x> = PolynomialRing(QQ)
sage: l = P.weil_polynomials(d=12, q=2)
sage: len(l)
164937
sage: l[0]
x^12 - 3*x^11 + 4*x^10 - 2*x^9 + 3*x^8 - 17*x^7 + 36*x^6 - 34*x^5 + 12*x^4
      - 16*x^3 + 64*x^2 - 96*x + 64
```

There are additional “power user” features that are not fully exposed in SageMath. See me if you need to use them.

Exhausting over Weil polynomials

SageMath includes code (by me) to exhaust over Weil polynomials satisfying various constraints. This was used to build the tables of isogeny classes of abelian varieties over finite fields in [LMFDB](#), but you can also use it directly.

```
sage: P.<x> = PolynomialRing(QQ)
sage: l = P.weil_polynomials(d=12, q=2)
sage: len(l)
164937
sage: l[0]
x^12 - 3*x^11 + 4*x^10 - 2*x^9 + 3*x^8 - 17*x^7 + 36*x^6 - 34*x^5 + 12*x^4
      - 16*x^3 + 64*x^2 - 96*x + 64
```

There are additional “power user” features that are not fully exposed in SageMath. See me if you need to use them.

Exhausting over Weil polynomials

SageMath includes code (by me) to exhaust over Weil polynomials satisfying various constraints. This was used to build the tables of isogeny classes of abelian varieties over finite fields in [LMFDB](#), but you can also use it directly.

```
sage: P.<x> = PolynomialRing(QQ)
sage: l = P.weil_polynomials(d=12, q=2)
sage: len(l)
164937
sage: l[0]
x^12 - 3*x^11 + 4*x^10 - 2*x^9 + 3*x^8 - 17*x^7 + 36*x^6 - 34*x^5 + 12*x^4
      - 16*x^3 + 64*x^2 - 96*x + 64
```

There are additional “power user” features that are not fully exposed in SageMath. See me if you need to use them.

Computing zeta functions

In principle, one can compute the zeta function of a K3 surface X over $\mathbb{F}_q$ by enumerating $X(\mathbb{F}_{q^n})$ for $n = 1, \dots, 10$ plus the sign of the functional equation.⁵ In practice this is only feasible for $q = 2$ or in cases with a big Picard sublattice (with known Galois action).

Instead, one can interpret $Q(T)$ as the (reversed) charpoly of Frobenius on H^2 for a suitable cohomology theory. The most practical for computation is **not** étale (ℓ -adic) cohomology but rather crystalline (p -adic) cohomology. This is a long story with contributions from many people (K, Tuitman, Harvey, Costa, etc.).

Executive summary: if you choose a smooth lift of X over the Witt vectors $W(\mathbb{F}_q)$, then the algebraic de Rham cohomology admits a Frobenius action given by a certain p -adically convergent power series. Expressing this in terms of generators of cohomology involves a lot of pole order reduction (e.g., Griffiths–Dwork for a projective hypersurface).

⁵Even without the sign, one has at most two candidates for $Q(T)$, and in many cases only one has roots on the unit circle.

Computing zeta functions

In principle, one can compute the zeta function of a K3 surface X over $\mathbb{F}_q$ by enumerating $X(\mathbb{F}_{q^n})$ for $n = 1, \dots, 10$ plus the sign of the functional equation.⁵ In practice this is only feasible for $q = 2$ or in cases with a big Picard sublattice (with known Galois action).

Instead, one can interpret $Q(T)$ as the (reversed) charpoly of Frobenius on H^2 for a suitable cohomology theory. The most practical for computation is **not** étale (ℓ -adic) cohomology but rather crystalline (p -adic) cohomology. This is a long story with contributions from many people (K, Tuitman, Harvey, Costa, etc.).

Executive summary: if you choose a smooth lift of X over the Witt vectors $W(\mathbb{F}_q)$, then the algebraic de Rham cohomology admits a Frobenius action given by a certain p -adically convergent power series. Expressing this in terms of generators of cohomology involves a lot of pole order reduction (e.g., Griffiths–Dwork for a projective hypersurface).

⁵Even without the sign, one has at most two candidates for $Q(T)$, and in many cases only one has roots on the unit circle.

Computing zeta functions

In principle, one can compute the zeta function of a K3 surface X over $\mathbb{F}_q$ by enumerating $X(\mathbb{F}_{q^n})$ for $n = 1, \dots, 10$ plus the sign of the functional equation.⁵ In practice this is only feasible for $q = 2$ or in cases with a big Picard sublattice (with known Galois action).

Instead, one can interpret $Q(T)$ as the (reversed) charpoly of Frobenius on H^2 for a suitable cohomology theory. The most practical for computation is **not** étale (ℓ -adic) cohomology but rather crystalline (p -adic) cohomology. This is a long story with contributions from many people (K, Tuitman, Harvey, Costa, etc.).

Executive summary: if you choose a smooth lift of X over the Witt vectors $W(\mathbb{F}_q)$, then the algebraic de Rham cohomology admits a Frobenius action given by a certain p -adically convergent power series. Expressing this in terms of generators of cohomology involves a lot of pole order reduction (e.g., Griffiths–Dwork for a projective hypersurface).

⁵Even without the sign, one has at most two candidates for $Q(T)$, and in many cases only one has roots on the unit circle.

Census data

Census data exists for K3 surfaces over $\mathbb{F}_2$ of degrees 2, 4, 6, 8 (by work of K–Sutherland, Auel–Frei–Mandel, and Auel–Shaikh). In each case, for all **smooth** members of the corresponding family,⁶ isomorphism class representatives and zeta functions (computed by enumeration) are available.

However, one would like this also for members of the family with ADE singularities. In particular this would add many examples with elevated geometric Picard rank, and in particular many more supersingular examples (perhaps with new values of Q_{alg}).

A related task is making censuses of curves of genus g over $\mathbb{F}_q$ (Huang–K–Lau). Recently Bergström has reported some success using AI to accelerate these censuses.

⁶Over a finite field, there are no Brauer–Severi varieties. So for example in degree 4 we really only need to consider quartics in $\mathbb{P}^3$.

Census data

Census data exists for K3 surfaces over $\mathbb{F}_2$ of degrees 2, 4, 6, 8 (by work of K–Sutherland, Auel–Frei–Mandel, and Auel–Shaikh). In each case, for all **smooth** members of the corresponding family,⁶ isomorphism class representatives and zeta functions (computed by enumeration) are available.

However, one would like this also for members of the family with ADE singularities. In particular this would add many examples with elevated geometric Picard rank, and in particular many more supersingular examples (perhaps with new values of Q_{alg}).

A related task is making censuses of curves of genus g over $\mathbb{F}_q$ (Huang–K–Lau). Recently Bergström has reported some success using AI to accelerate these censuses.

⁶Over a finite field, there are no Brauer–Severi varieties. So for example in degree 4 we really only need to consider quartics in $\mathbb{P}^3$.

Census data

Census data exists for K3 surfaces over $\mathbb{F}_2$ of degrees 2, 4, 6, 8 (by work of K–Sutherland, Auel–Frei–Mandel, and Auel–Shaikh). In each case, for all **smooth** members of the corresponding family,⁶ isomorphism class representatives and zeta functions (computed by enumeration) are available.

However, one would like this also for members of the family with ADE singularities. In particular this would add many examples with elevated geometric Picard rank, and in particular many more supersingular examples (perhaps with new values of Q_{alg}).

A related task is making censuses of curves of genus g over $\mathbb{F}_q$ (Huang–K–Lau). Recently Bergström has reported some success using AI to accelerate these censuses.

⁶Over a finite field, there are no Brauer–Severi varieties. So for example in degree 4 we really only need to consider quartics in $\mathbb{P}^3$.

Contents

- 1 Prelude on zeta functions
- 2 Structure of the zeta function of a K3 surface
- 3 Positivity considerations
- 4 Computational aspects
- 5 Lattice-theoretic considerations**
- 6 Conclusion

The Galois action on Néron–Severi

For X a K3 surface over $\mathbb{F}_q$, the group $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is finite free of rank $\bar{\rho} \leq 22$ and carries the intersection pairing. By the Hodge index theorem, the pairing has signature $(1, \bar{\rho} - 1)$.

The absolute Galois group $G_{\mathbb{F}_q}$ acts on $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ preserving the pairing and the ample cone. For $\text{Frob}_q \in G_{\mathbb{F}_q}$ the canonical generator, we have

$$Q_{\text{alg}}(T) = \det(1 - \text{Frob}_q T, \text{Pic}(X_{\overline{\mathbb{F}}_q})).$$

The map $\text{Pic}(X) \rightarrow \text{Pic}(X_{\overline{\mathbb{F}}_q})$ is a primitive embedding with image $\text{Pic}(X_{\overline{\mathbb{F}}_q})^{G_{\mathbb{F}_q}}$.

The Galois action on Néron–Severi

For X a K3 surface over $\mathbb{F}_q$, the group $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is finite free of rank $\bar{\rho} \leq 22$ and carries the intersection pairing. By the Hodge index theorem, the pairing has signature $(1, \bar{\rho} - 1)$.

The absolute Galois group $G_{\mathbb{F}_q}$ acts on $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ preserving the pairing and the ample cone. For $\text{Frob}_q \in G_{\mathbb{F}_q}$ the canonical generator, we have

$$Q_{\text{alg}}(T) = \det(1 - \text{Frob}_q T, \text{Pic}(X_{\overline{\mathbb{F}}_q})).$$

The map $\text{Pic}(X) \rightarrow \text{Pic}(X_{\overline{\mathbb{F}}_q})$ is a primitive embedding with image $\text{Pic}(X_{\overline{\mathbb{F}}_q})^{G_{\mathbb{F}_q}}$.

The supersingular case

We have $\bar{\rho} = 22$ iff X is supersingular. By Rudakov–Sharafevich, the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is uniquely determined by p and the **Artin invariant** $\sigma \in \{1, \dots, 10\}$, which is characterized by

$$A_{\text{Pic}(X_{\overline{\mathbb{F}}_q})} \cong p^{2\sigma}.$$

The crystalline Torelli theorem was proved by Ogus for q odd:⁷ an isomorphism $X \rightarrow Y$ of supersingular K3 surfaces over $\mathbb{F}_q$ is specified by an isomorphism $\text{Pic}(X_{\overline{\mathbb{F}}_q}) \rightarrow \text{Pic}(Y_{\overline{\mathbb{F}}_q})$ preserving all of

- the Galois action;
- the intersection pairing;
- the ample cone;
- the kernel of the crystalline cycle class map.

⁷It may be possible to extend this to even q by replacing crystalline cohomology with **prismatic cohomology** (Bhatt–Scholze) and the corresponding cycle class map.

The supersingular case

We have $\bar{\rho} = 22$ iff X is supersingular. By Rudakov–Sharafevich, the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is uniquely determined by p and the **Artin invariant** $\sigma \in \{1, \dots, 10\}$, which is characterized by

$$A_{\text{Pic}(X_{\overline{\mathbb{F}}_q})} \cong p^{2\sigma}.$$

The crystalline Torelli theorem was proved by Ogus for q odd:⁷ an isomorphism $X \rightarrow Y$ of supersingular K3 surfaces over $\mathbb{F}_q$ is specified by an isomorphism $\text{Pic}(X_{\overline{\mathbb{F}}_q}) \rightarrow \text{Pic}(Y_{\overline{\mathbb{F}}_q})$ preserving all of

- the Galois action;
- the intersection pairing;
- the ample cone;
- the kernel of the crystalline cycle class map.

⁷It may be possible to extend this to even q by replacing crystalline cohomology with **prismatic cohomology** (Bhatt–Scholze) and the corresponding cycle class map.

The supersingular case

We have $\bar{p} = 22$ iff X is supersingular. By Rudakov–Sharafevich, the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is uniquely determined by p and the **Artin invariant** $\sigma \in \{1, \dots, 10\}$, which is characterized by

$$A_{\text{Pic}(X_{\overline{\mathbb{F}}_q})} \cong p^{2\sigma}.$$

The crystalline Torelli theorem was proved by Ogus for q odd:⁷ an isomorphism $X \rightarrow Y$ of supersingular K3 surfaces over $\mathbb{F}_q$ is specified by an isomorphism $\text{Pic}(X_{\overline{\mathbb{F}}_q}) \rightarrow \text{Pic}(Y_{\overline{\mathbb{F}}_q})$ preserving all of

- the Galois action;
- the intersection pairing;
- the ample cone;
- the kernel of the crystalline cycle class map.

⁷It may be possible to extend this to even q by replacing crystalline cohomology with **prismatic cohomology** (Bhatt–Scholze) and the corresponding cycle class map.

The supersingular case

We have $\bar{\rho} = 22$ iff X is supersingular. By Rudakov–Sharafevich, the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is uniquely determined by p and the **Artin invariant** $\sigma \in \{1, \dots, 10\}$, which is characterized by

$$A_{\text{Pic}(X_{\overline{\mathbb{F}}_q})} \cong p^{2\sigma}.$$

The crystalline Torelli theorem was proved by Ogus for q odd:⁷ an isomorphism $X \rightarrow Y$ of supersingular K3 surfaces over $\mathbb{F}_q$ is specified by an isomorphism $\text{Pic}(X_{\overline{\mathbb{F}}_q}) \rightarrow \text{Pic}(Y_{\overline{\mathbb{F}}_q})$ preserving all of

- the Galois action;
- the intersection pairing;
- the ample cone;
- the kernel of the crystalline cycle class map.

⁷It may be possible to extend this to even q by replacing crystalline cohomology with **prismatic cohomology** (Bhatt–Scholze) and the corresponding cycle class map.

The supersingular case

We have $\bar{\rho} = 22$ iff X is supersingular. By Rudakov–Sharafevich, the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is uniquely determined by p and the **Artin invariant** $\sigma \in \{1, \dots, 10\}$, which is characterized by

$$A_{\text{Pic}(X_{\overline{\mathbb{F}}_q})} \cong p^{2\sigma}.$$

The crystalline Torelli theorem was proved by Ogus for q odd:⁷ an isomorphism $X \rightarrow Y$ of supersingular K3 surfaces over $\mathbb{F}_q$ is specified by an isomorphism $\text{Pic}(X_{\overline{\mathbb{F}}_q}) \rightarrow \text{Pic}(Y_{\overline{\mathbb{F}}_q})$ preserving all of

- the Galois action;
- the intersection pairing;
- the ample cone;
- the kernel of the crystalline cycle class map.

⁷It may be possible to extend this to even q by replacing crystalline cohomology with **prismatic cohomology** (Bhatt–Scholze) and the corresponding cycle class map.

The supersingular case

We have $\bar{\rho} = 22$ iff X is supersingular. By Rudakov–Sharafevich, the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is uniquely determined by p and the **Artin invariant** $\sigma \in \{1, \dots, 10\}$, which is characterized by

$$A_{\text{Pic}(X_{\overline{\mathbb{F}}_q})} \cong p^{2\sigma}.$$

The crystalline Torelli theorem was proved by Ogus for q odd:⁷ an isomorphism $X \rightarrow Y$ of supersingular K3 surfaces over $\mathbb{F}_q$ is specified by an isomorphism $\text{Pic}(X_{\overline{\mathbb{F}}_q}) \rightarrow \text{Pic}(Y_{\overline{\mathbb{F}}_q})$ preserving all of

- the Galois action;
- the intersection pairing;
- the ample cone;
- the kernel of the crystalline cycle class map.

⁷It may be possible to extend this to even q by replacing crystalline cohomology with **prismatic cohomology** (Bhatt–Scholze) and the corresponding cycle class map.

The supersingular case

We have $\bar{\rho} = 22$ iff X is supersingular. By Rudakov–Sharafevich, the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ is uniquely determined by p and the **Artin invariant** $\sigma \in \{1, \dots, 10\}$, which is characterized by

$$A_{\text{Pic}(X_{\overline{\mathbb{F}}_q})} \cong p^{2\sigma}.$$

The crystalline Torelli theorem was proved by Ogus for q odd:⁷ an isomorphism $X \rightarrow Y$ of supersingular K3 surfaces over $\mathbb{F}_q$ is specified by an isomorphism $\text{Pic}(X_{\overline{\mathbb{F}}_q}) \rightarrow \text{Pic}(Y_{\overline{\mathbb{F}}_q})$ preserving all of

- the Galois action;
- the intersection pairing;
- the ample cone;
- the kernel of the crystalline cycle class map.

⁷It may be possible to extend this to even q by replacing crystalline cohomology with **prismatic cohomology** (Bhatt–Scholze) and the corresponding cycle class map.

Artin–Tate formula

The Artin–Tate formula (true because of the Tate conjecture!) asserts that

$$\lim_{T \rightarrow 1} \frac{Q(T)}{(T-1)^{\text{ord}_{T=1}(Q)}} = \#A_{\text{Pic}(X)} \# \text{Br}(X).$$

The group $\# \text{Br}(X)$ carries an alternating pairing, so its order is a square.

The same formula applies after base extension. Using this we can write down candidates for the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ and its Galois action that are consistent with Artin–Tate.

Remark of Elsenhans–Jahnel: sometimes there are no candidates! When $\text{ord}_{T=-1}(Q) = 0$, the map $\text{Pic}(X) \rightarrow \text{Pic}(X_{\mathbb{F}_{q^2}})$ is an isomorphism and so $\#A_{\text{Pic}(X)} = \#A_{\text{Pic}(X_{\mathbb{F}_{q^2}})}$; since $\text{ord}_{T=1}(Q) + \text{ord}_{T=-1}(Q)$ is always even, this implies that $Q(-1)$ must be a square.

Artin–Tate formula

The Artin–Tate formula (true because of the Tate conjecture!) asserts that

$$\lim_{T \rightarrow 1} \frac{Q(T)}{(T-1)^{\text{ord}_{T=1}(Q)}} = \#A_{\text{Pic}(X)} \# \text{Br}(X).$$

The group $\# \text{Br}(X)$ carries an alternating pairing, so its order is a square.

The same formula applies after base extension. Using this we can write down candidates for the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ and its Galois action that are consistent with Artin–Tate.

Remark of Elsenhans–Jahnel: sometimes there are no candidates! When $\text{ord}_{T=-1}(Q) = 0$, the map $\text{Pic}(X) \rightarrow \text{Pic}(X_{\mathbb{F}_{q^2}})$ is an isomorphism and so $\#A_{\text{Pic}(X)} = \#A_{\text{Pic}(X_{\mathbb{F}_{q^2}})}$; since $\text{ord}_{T=1}(Q) + \text{ord}_{T=-1}(Q)$ is always even, this implies that $Q(-1)$ must be a square.

Artin–Tate formula

The Artin–Tate formula (true because of the Tate conjecture!) asserts that

$$\lim_{T \rightarrow 1} \frac{Q(T)}{(T-1)^{\text{ord}_{T=1}(Q)}} = \#A_{\text{Pic}(X)} \# \text{Br}(X).$$

The group $\# \text{Br}(X)$ carries an alternating pairing, so its order is a square.

The same formula applies after base extension. Using this we can write down candidates for the lattice $\text{Pic}(X_{\overline{\mathbb{F}}_q})$ and its Galois action that are consistent with Artin–Tate.

Remark of Elsenhans–Jahnel: sometimes there are no candidates! When $\text{ord}_{T=-1}(Q) = 0$, the map $\text{Pic}(X) \rightarrow \text{Pic}(X_{\mathbb{F}_{q^2}})$ is an isomorphism and so $\#A_{\text{Pic}(X)} = \#A_{\text{Pic}(X_{\mathbb{F}_{q^2}})}$; since $\text{ord}_{T=1}(Q) + \text{ord}_{T=-1}(Q)$ is always even, this implies that $Q(-1)$ must be a square.

Lifts to characteristic zero

Now suppose that X is not supersingular, so $\bar{\rho} \leq 20$. In this case, there always exists a lift $\mathfrak{X}$ of X over some finite totally ramified extension $\mathfrak{o}$ of $W(\mathbb{F}_q)$ such that the reduction map

$$\mathrm{Pic}(\mathfrak{X}_{\mathbb{C}}) \cong \mathrm{Pic}(\mathfrak{X}) \rightarrow \mathrm{Pic}(X_{\overline{\mathbb{F}}_q})$$

is an isomorphism (normally it is only injective).

This means that just as over $\mathbb{C}$, $\mathrm{Pic}(X_{\overline{\mathbb{F}}_q})$ (which has signature $(1, \bar{\rho} - 1)$) admits a primitive embedding into the K3 lattice $E_8^{\oplus 2} \oplus U^{\oplus 3}$. There is no way to construct this directly from X because there is no meaningful interpretation of $H^2(X_{\overline{\mathbb{F}}_q}, \mathbb{Z})$.

For $p > 2$, Inoue⁸ established the existence of a **quasi-canonical lift** of X . In particular the transcendental Hodge structure of the lift has CM in the sense of Zarhin.

⁸For $p > 3$ this was done by Nygaard–Ogus. For $p = 2$, Ito–Ito–Koshikawa produce a quasi-canonical lift after a base extension.

Lifts to characteristic zero

Now suppose that X is not supersingular, so $\bar{\rho} \leq 20$. In this case, there always exists a lift $\mathfrak{X}$ of X over some finite totally ramified extension $\mathfrak{o}$ of $W(\mathbb{F}_q)$ such that the reduction map

$$\mathrm{Pic}(\mathfrak{X}_{\mathbb{C}}) \cong \mathrm{Pic}(\mathfrak{X}) \rightarrow \mathrm{Pic}(X_{\overline{\mathbb{F}}_q})$$

is an isomorphism (normally it is only injective).

This means that just as over $\mathbb{C}$, $\mathrm{Pic}(X_{\overline{\mathbb{F}}_q})$ (which has signature $(1, \bar{\rho} - 1)$) admits a primitive embedding into the K3 lattice $E_8^{\oplus 2} \oplus U^{\oplus 3}$. There is no way to construct this directly from X because there is no meaningful interpretation of $H^2(X_{\overline{\mathbb{F}}_q}, \mathbb{Z})$.

For $p > 2$, Inoue⁸ established the existence of a **quasi-canonical lift** of X . In particular the transcendental Hodge structure of the lift has CM in the sense of Zarhin.

⁸For $p > 3$ this was done by Nygaard–Ogus. For $p = 2$, Ito–Ito–Koshikawa produce a quasi-canonical lift after a base extension.

Lifts to characteristic zero

Now suppose that X is not supersingular, so $\bar{\rho} \leq 20$. In this case, there always exists a lift $\mathfrak{X}$ of X over some finite totally ramified extension $\mathfrak{o}$ of $W(\mathbb{F}_q)$ such that the reduction map

$$\mathrm{Pic}(\mathfrak{X}_{\mathbb{C}}) \cong \mathrm{Pic}(\mathfrak{X}) \rightarrow \mathrm{Pic}(X_{\overline{\mathbb{F}}_q})$$

is an isomorphism (normally it is only injective).

This means that just as over $\mathbb{C}$, $\mathrm{Pic}(X_{\overline{\mathbb{F}}_q})$ (which has signature $(1, \bar{\rho} - 1)$) admits a primitive embedding into the K3 lattice $E_8^{\oplus 2} \oplus U^{\oplus 3}$. There is no way to construct this directly from X because there is no meaningful interpretation of $H^2(X_{\overline{\mathbb{F}}_q}, \mathbb{Z})$.

For $p > 2$, Inoue⁸ established the existence of a **quasi-canonical lift** of X . In particular the transcendental Hodge structure of the lift has CM in the sense of Zarhin.

⁸For $p > 3$ this was done by Nygaard–Ogus. For $p = 2$, Ito–Ito–Koshikawa produce a quasi-canonical lift after a base extension.

Dare to dream: Deligne modules?

One can use CM lifts to give “linear algebraic” descriptions of the category of abelian varieties over $\mathbb{F}_q$ (Deligne, Centeleghe–Stix, Bergström–Karemaker–Marseglia, Rybakov). These are practical for tabulating **isomorphism classes** of abelian varieties (Marseglia), as opposed to **isogeny classes** which currently occur in LMFDB.

Concrete example: define a **Deligne module** to be a finite free abelian group with two operators F, V such that $FV = VF = q$ and $F \otimes \mathbb{Q}$ is semisimple. Deligne showed that the category of **ordinary** abelian varieties over $\mathbb{F}_q$ is equivalent to the category of Deligne modules for which the characteristic polynomial of $F \otimes \mathbb{Q}$ is an ordinary q -Weil polynomial.

Question

Is there a corresponding description of the category of K3 surfaces over $\mathbb{F}_q$?

This would potentially help us answer many of the questions raised earlier...

Dare to dream: Deligne modules?

One can use CM lifts to give “linear algebraic” descriptions of the category of abelian varieties over $\mathbb{F}_q$ (Deligne, Centeleghe–Stix, Bergström–Karemaker–Marseglia, Rybakov). These are practical for tabulating **isomorphism classes** of abelian varieties (Marseglia), as opposed to **isogeny classes** which currently occur in LMFDB.

Concrete example: define a **Deligne module** to be a finite free abelian group with two operators F, V such that $FV = VF = q$ and $F \otimes \mathbb{Q}$ is semisimple. Deligne showed that the category of **ordinary** abelian varieties over $\mathbb{F}_q$ is equivalent to the category of Deligne modules for which the characteristic polynomial of $F \otimes \mathbb{Q}$ is an ordinary q -Weil polynomial.

Question

Is there a corresponding description of the category of K3 surfaces over $\mathbb{F}_q$?

This would potentially help us answer many of the questions raised earlier...

Dare to dream: Deligne modules?

One can use CM lifts to give “linear algebraic” descriptions of the category of abelian varieties over $\mathbb{F}_q$ (Deligne, Centeleghe–Stix, Bergström–Karemaker–Marseglia, Rybakov). These are practical for tabulating **isomorphism classes** of abelian varieties (Marseglia), as opposed to **isogeny classes** which currently occur in LMFDB.

Concrete example: define a **Deligne module** to be a finite free abelian group with two operators F, V such that $FV = VF = q$ and $F \otimes \mathbb{Q}$ is semisimple. Deligne showed that the category of **ordinary** abelian varieties over $\mathbb{F}_q$ is equivalent to the category of Deligne modules for which the characteristic polynomial of $F \otimes \mathbb{Q}$ is an ordinary q -Weil polynomial.

Question

Is there a corresponding description of the category of K3 surfaces over $\mathbb{F}_q$?

This would potentially help us answer many of the questions raised earlier...

Dare to dream: Deligne modules?

One can use CM lifts to give “linear algebraic” descriptions of the category of abelian varieties over $\mathbb{F}_q$ (Deligne, Centeleghe–Stix, Bergström–Karemaker–Marseglia, Rybakov). These are practical for tabulating **isomorphism classes** of abelian varieties (Marseglia), as opposed to **isogeny classes** which currently occur in LMFDB.

Concrete example: define a **Deligne module** to be a finite free abelian group with two operators F, V such that $FV = VF = q$ and $F \otimes \mathbb{Q}$ is semisimple. Deligne showed that the category of **ordinary** abelian varieties over $\mathbb{F}_q$ is equivalent to the category of Deligne modules for which the characteristic polynomial of $F \otimes \mathbb{Q}$ is an ordinary q -Weil polynomial.

Question

Is there a corresponding description of the category of K3 surfaces over $\mathbb{F}_q$?

This would potentially help us answer many of the questions raised earlier...

Contents

- 1 Prelude on zeta functions
- 2 Structure of the zeta function of a K3 surface
- 3 Positivity considerations
- 4 Computational aspects
- 5 Lattice-theoretic considerations
- 6 Conclusion

Selected references

- M. Artin, Supersingular $K3$ surfaces, *Ann. Scient. Éc. Norm. Sup.* **4** (1974), 543—568.
- F. Bernasconi and S. Filipazzi, Rational points on 3-folds with nef anti-canonical class over finite fields, *Proc. London Math. Soc.* **130** (2025).
- T.G. Centeleghe and J. Stix, Categories of abelian varieties over finite fields, I: Abelian varieties over $\mathbb{F}_p$, *Algebra and Number Theory* **9** (2015), 225–265.
- P. Deligne, Variétés abéliennes ordinaires sur un corps fini, *Invent. Math.* **8** (1969), 238—243.
- K. Inoue, Construction of quasi-canonical liftings of $K3$ surfaces of finite height in odd characteristic, *manuscripta math.* **173** (2024), 619–635.
- K. Ito, Unconditional construction of $K3$ surfaces over finite fields with given L -function in large characteristic, *manuscripta math.* **159** (2019), 281–300.
- K. Ito, T. Ito, and T. Koshikawa, CM liftings of $K3$ surfaces over finite fields and their applications to the Tate conjecture, *Forum of Math. Sigma* **9** (2021), 1–70.
- K.S. Kedlaya, Remarks on the zeta function of a $K3$ surface over a finite field, [preprint](#).
- J.S. Milne and W.C. Waterhouse, Abelian varieties over finite fields, *Number Theory Institute (State Univ. New York, Stony Brook, N.Y., 1969)*, Proc. Sympos. Pure Math. 20, Amer. Math. Soc., Providence, 1971, 53—64.
- N. Nygaard and A. Ogus, Tate’s conjecture for $K3$ surfaces of finite height, *Annals of Math.* **122** (1985), 461–507.
- A. Ogus, A crystalline Torelli theorem for supersingular $K3$ surfaces, in *Arithmetic and Geometry*, Progress in Mathematics 36, Birkhäuser, Boston, MA, 1983.

That QR code again

